

(FILE 'HOME' ENTERED AT 10:37:11 ON 17 MAR 2004)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, CANCERLIT, JAPIO' ENTERED AT  
10:37:29 ON 17 MAR 2004

L1	21773 S NEURITE? AND OUTGROWTH?
L2	40 S L1 AND LUMINES?
L3	391 S NEUROFILAMENT AND IMAGE?
L4	1 S L2 AND L3
L5	25 S L3 AND L1
L6	12 DUPLICATE REMOVE L5 (13 DUPLICATES REMOVED)
L7	11 S L6 NOT L4
L8	2359 S L1 AND NUCLE?
L9	2 S L8 AND L3

(FILE 'HOME' ENTERED AT 10:37:11 ON 17 MAR 2004)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, CANCERLIT, JAPIO' ENTERED AT  
10:37:29 ON 17 MAR 2004

L1	21773 S NEURITE? AND OUTGROWTH?
L2	40 S L1 AND LUMINES?
L3	391 S NEUROFILAMENT AND IMAGE?
L4	1 S L2 AND L3
L5	25 S L3 AND L1
L6	12 DUPLICATE REMOVE L5 (13 DUPLICATES REMOVED)
L7	11 S L6 NOT L4
L8	2359 S L1 AND NUCLE?
L9	2 S L8 AND L3



NPL Virtual Library

My List - 0 Help

Search

[Main Search](#) | [Advanced Keyword Search](#) | [Search History](#)

Search: ISSN

0305-1870

[Refine Search](#)

You're searching: Scientific and Technical Information Center

## Item Information

- Subscriptions
- Holdings

## Browse Catalog

by title:

- Clinical and experim...

## MARC Display

### Clinical and experimental pharmacology and physiology [electronic resource].

Imprint: Oxford, England : Blackwell Pub. Ltd.,

URL: <http://ejournals.ebsco.com/Journal.asp?JournalID=101685> Click here to see page images available via Ebsco. v.26, n.1 (1999)-  
<http://search.epnet.com/direct.asp?db=aph&jid=%22ETO%22&scope=site> Click here for Online version via Academic Search Premium (ASP). Jan 1998-

Notes: Available on ADONIS, v. 27, no. 1-2 (2000) - v. 29, S4 (2002)  
 Mode of access: World Wide Web.

ISSN: 1440-1681  
 0305-1870 (Print ed.)

Subjects: Clinical pharmacology -- Periodicals.  
 Pharmacology, Experimental -- Periodicals.  
 Physiology, Pathological -- Periodicals.  
 Physiology, Experimental -- Periodicals.

## Subscription Summary

## US Patent &amp; Trademark Office

Location: US Patent & Trademark Office  
 Collection: Electronic  
 Call No.: e-journal (Ebsco)  
 Status: Currently Received  
 Media Type: Elec

[Show all items](#)

Email: [katherine.arendt@uspto.gov](mailto:katherine.arendt@uspto.gov) to ask questions or make suggestions.

iPac 2.03.01

Brought to you by *Scientific and Technical Information Center*

AN 1999:198128 CAPLUS  
 DN 131:16923  
 ED Entered STN: 29 Mar 1999  
 TI Effects of different fragments of tenascin-R on neuron morphology in vitro  
 AU Xu, Hanpeng; Xiao, Huasheng; Wang, Haojun; Liang, Zhe; Gong, Ju  
 CS The Institute of Neuroscience, Fourth Military Medical University, Xi'an, 710033, Peop. Rep. China  
 SO Journal of Medical Colleges of PLA (1998), 13(4), 272-275  
 CODEN: JMCPE6; ISSN: 1000-1948  
 PB Journal of Medical Colleges of PLA, Editorial Board  
 DT Journal  
 LA English  
 CC 13-6 (Mammalian Biochemistry)  
 AB To investigate the effects of different tenascin-R fragments on morphol. changes of neurons in vitro, cell suspension were prepared from E14-15 mouse embryo spinal cords by mech. dissection and trypsin digestion. The cells were cultured in dishes coated with different bacterial expressed tenascin-R fragments. After being cultured in serum-free medium for 26 h, the cells were fixed and stained by ABC immunocytochem. method for NSE. The cell number and **neurite** length were measured by a stereol. method using an **image** anal. system, and the data was analyzed statistically. The cells grew well in the serum-free medium for 26 h. Three types of cells were identified: (1) phase-bright cells with single or double **neurites**; (2) phase-dark cells with well branching **neurites**; (3) flat cells with 2-4 round vesicles in the **cell body** and radio-like **neurites**. The cell number and the **neurites** length were influenced by different tenascin-R fragments. It was found that FN1-2 fragment inhibited **neurite outgrowth**. Different tenascin-R fragments that were used as substrate exert varying effects on cultured neural cells, adhesion or anti-adhesion of cells, promotion or inhibition of the growth of **neurites**. These influences were mediated through receptors on the cell membrane. This study may provide some clues to the search for the search for the different receptors which may play essential roles in neuronal development and plasticity.  
 ST tenascin R neuron morphol  
 IT Nerve  
 (neuron; tenascin-R fragment effect on neuron)  
 IT Axon  
 (outgrowth; tenascin-R fragment effect on neuron)  
 IT Cell adhesion  
 Cell morphology  
 Spinal cord  
 (tenascin-R fragment effect on neuron)  
 IT Tenascins  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
 (tenascin-R; tenascin-R fragment effect on neuron)  
 RE.CNT 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 RE  
 (1) Chiquet, E; Curr Opin Cell Biol 1995, V7(5), P715  
 (2) Dorries, U; J Neurosci Res 1996, V43(4), P420 MEDLINE  
 (3) Gundersen, H; APMIS 1988, V96(5), P379 MEDLINE  
 (4) Gundersen, H; J Microsc 1987, V147(Pt3), P229  
 (5) Hay Elizabeth, D; Cell Biol of ECM 2nd ed 1991, P419  
 (6) Kruse, J; Nature 1985, V316, P146 CAPLUS  
 (7) Pesheva, P; J Cell Biol 1989, V109(4), P1765 CAPLUS  
 (8) Pesheva, P; Neuron 1992, V6(1), P69  
 (9) Sjmith, B; Cell Adv Com 1996, V3(6), P451  
 (10) Taylor, J; J Neurosci Res 1993, V35(4), P347 CAPLUS  
 (11) Vidal, L; J Neurosci Methods 1995, V56(1), P89 CAPLUS

AN 1998:517015 CAPLUS  
 DN 129:273372  
 ED Entered STN: 20 Aug 1998  
 TI p21RhoA and p21RhoA binding proteins as regulators of lysophosphatidic acid (LPA)-induced changes in neuronal morphology  
 AU Gibbink, Martijn F. B. G.; Kranenburg, Onno; Jalink, Kees; Postma, Friso R.; Poland, Mieke; Houssa, Brahim; Oomen, Lauren; Van Horck, Francis P. G.; Moolenaar, Wouter H.  
 CS Divison Cellular Biochemistry, The Netherlands Cancer Institute, Amsterdam, 1066 CX, Neth.  
 SO Kinases and Phosphatases in Lymphocyte and Neuronal Signaling (1997), 235-241. Editor(s): Yakura, Hidetaka. Publisher: Springer, Tokyo, Japan. CODEN: 66NHAY  
 DT Conference; General Review  
 LA English  
 CC 13-0 (Mammalian Biochemistry)  
 AB This is a **review** with 34 refs. Addition of lysophosphatidic acid (LPA), sphingosine-1-phosphate or thrombin to serum-deprived N1E-15 neuronal cells results in rapid neurite retraction and rounding of the **cell body**. These morphol. changes are accompanied by rapid assembly of filamentous actin and contraction of the cortical actin cytoskeleton. Treatment of the cells with Clostridium botulinum C3 exoenzyme, which ADP-ribosylates and thereby inactivates the Rho small GTP binding proteins, inhibits agonist induced cell rounding. Furthermore, dominant neg. RhoA stimulates cell flattening and **neurite outgrowth** similar to C3 toxin. Cells expressing dominant-neg. RhoA also fail to change shape in response to LPA. Activated V14RhoA mimics LPA action in inducing cell rounding and inhibiting neuronal outgrowth. To further elucidate the signaling mechanisms that regulate RhoA mediated changes in neuronal morphol., novel RhoA binding proteins were identified. RhoGEF is a novel GDP/GTP exchanger that interacts with both wild-type and activated V14RhoA, but not with Rac or CDC42. Similar to activated V14RhoA, RhoGEF induces cell rounding. p116Rip is a novel RhoA binding protein whose overexpression stimulates **neurite outgrowth**. These results establish RhoA, RhoGEF and p116Rip as critical determinants of RhoA-mediated neuronal shape changes. Identification of proteins that regulate LPA induced actin cytoskeletal remodeling will help to understand to intracellular mechanisms by which neuronal cells regulate their complex morphol.  
 ST **review** p21RhoA protein neuronal morphol regulator; p116Rip  
 RhoGEF protein neuronal morphol **review**  
 IT Proteins, specific or class  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
 (RhoGEF; p21RhoA and p21RhoA binding proteins as regulators of lysophosphatidic acid (LPA)-induced changes in neuronal morphol.)  
 IT Nerve  
 (neuron; p21RhoA and p21RhoA binding proteins as regulators of lysophosphatidic acid (LPA)-induced changes in neuronal morphol.)  
 IT Proteins, specific or class  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
 (p116Rip; p21RhoA and p21RhoA binding proteins as regulators of lysophosphatidic acid (LPA)-induced changes in neuronal morphol.)  
 IT Cell morphology  
 Cytoskeleton  
 (p21RhoA and p21RhoA binding proteins as regulators of lysophosphatidic acid (LPA)-induced changes in neuronal morphol.)  
 IT Lysophosphatidic acids  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (p21RhoA and p21RhoA binding proteins as regulators of lysophosphatidic

acid (LPA)-induced changes in neuronal morphol.)

IT Rho protein (G protein)

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)

(p21rhoA; p21RhoA and p21RhoA binding proteins as regulators of lysophosphatidic acid (LPA)-induced changes in neuronal morphol.)

RE.CNT 34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Adra, C; Proc Natl Acad Sci U S A 1997, V94, P4279 CAPLUS
- (2) Amano, M; Science 1996, V271, P648 CAPLUS
- (3) Bunemann, M; EMBO J 1996, V15, P5527 MEDLINE
- (4) Fourcade, O; Cell 1995, V80, P919 CAPLUS
- (5) Gebbink, M; J Cell Biol in press 1997
- (6) Hecht, J; J Cell Biol 1996, V135, P1071 CAPLUS
- (7) Hotta, K; Biochem Biophys Res Commun 1996, V225, P69 CAPLUS
- (8) Ishizaki, T; EMBO J 1996, V15, P1885 CAPLUS
- (9) Jalink, K; Bioch Biophys Acta 1994, V1198, P185 CAPLUS
- (10) Jalink, K; Cell Growth Differ 1993, V4, P247 CAPLUS
- (11) Jalink, K; J Cell Biol 1992, V118, P411 CAPLUS
- (12) Jalink, K; J Cell Biol 1994, V126, P801 CAPLUS
- (13) Kamata, Y; Microbiol Immunol 1994, V38, P421 CAPLUS
- (14) Kimura, K; Science 1996, V273, P245 CAPLUS
- (15) Kranenburg, O; Submitted 1997
- (16) Machesky, L; Trends Cell Biol 1996, V6, P304 CAPLUS
- (17) Macrae, A; Mol Brain Res 1996, V42, P245 CAPLUS
- (18) Madaule, P; FEBS Lett 1995, V377, P243 CAPLUS
- (19) Matsui, T; EMBO J 1996, V15, P2208 CAPLUS
- (20) Moolenaar, W; Curr Opin Cell Biol 1995, V7, P203 CAPLUS
- (21) Moolenaar, W; Curr Opin Cell Biol 1997, V9, P168 CAPLUS
- (22) Moolenaar, W; J Biol Chem 1995, V270, P12949 CAPLUS
- (23) Ozaki, K; EMBO J 1996, V15, P2196 CAPLUS
- (24) Postma, F; EMBO J 1996, P2388 CAPLUS
- (25) Reid, T; J Biol Chem 1996, V271, P13556 CAPLUS
- (26) Ridley, A; Curr Biol 1996, V5, P1256
- (27) Schulze, C; J Neurochem 1997, V68, P991 CAPLUS
- (28) Symons, M; Trends Biochem Sci 1996, V21, P178 CAPLUS
- (29) Tigyi, G; J Neurochem 1996, V66, P537 CAPLUS
- (30) Toksoz, D; Oncogene 1995, V9, P621
- (31) van Koppen, C; J Biol Chem 1996, V271, P2082 CAPLUS
- (32) Whitehead, I; J Biol Chem 1995, V270, P18388 CAPLUS
- (33) Zalcman, G; J Biol Chem 1996, V271, P30366 CAPLUS
- (34) Zheng, Y; J Biol Chem 1995, V270, P9031 CAPLUS

d his

(FILE 'HOME' ENTERED AT 11:17:43 ON 17 MAR 2004)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, CANCERLIT, JAPIO' ENTERED AT  
11:18:09 ON 17 MAR 2004

L1 3714 S (CELL BOD?) AND NEURITE?  
L2 538 S L1 AND NUCLE?  
L3 113 S L2 AND OUTGROWTH?  
L4 4 S L3 AND IMAGE?  
L5 1 DUPLICATE REMOVE L4 (3 DUPLICATES REMOVED)  
L6 710 S (NEURITE OUTGROWTH) AND REVIEW  
L7 0 S L6 AND IMAGE?  
L8 8 S L6 AND (CELL BOD?)  
L9 7 DUPLICATE REMOVE L8 (1 DUPLICATE REMOVED)  
L10 38031 S FLUORES? AND LUMINES?  
L11 1055 S L10 AND REVIEW?  
L12 30 S L11 AND IMAGE?  
L13 25 DUPLICATE REMOVE L12 (5 DUPLICATES REMOVED)  
L14 0 S (FLUORE? VERSES LUMINES?)

=>



d his

(FILE 'HOME' ENTERED AT 11:17:43 ON 17 MAR 2004)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, CANCERLIT, JAPIO' ENTERED AT  
11:18:09 ON 17 MAR 2004

L1 3714 S (CELL BOD?) AND NEURITE?  
L2 538 S L1 AND NUCLE?  
L3 113 S L2 AND OUTGROWTH?  
L4 4 S L3 AND IMAGE?  
L5 1 DUPLICATE REMOVE L4 (3 DUPLICATES REMOVED)  
L6 710 S (NEURITE OUTGROWTH) AND REVIEW  
L7 0 S L6 AND IMAGE?  
L8 8 S L6 AND (CELL BOD?)  
L9 7 DUPLICATE REMOVE L8 (1 DUPLICATE REMOVED)  
L10 38031 S FLUORES? AND LUMINES?  
L11 1055 S L10 AND REVIEW?  
L12 30 S L11 AND IMAGE?  
L13 25 DUPLICATE REMOVE L12 (5 DUPLICATES REMOVED)  
L14 0 S (FLUORE? VERSES LUMINES?)

=>



SCIENCE @ DIRECT

Register or Login: Password: 
[Home](#) [Journals](#) [Abstract Databases](#) [Books](#) [Reference Works](#) [My Profile](#) [Alerts](#) [Help](#)
WELCOME GUEST USER [Info](#)**Developmental Brain Research**

Volume 108, Issues 1-2, 15 June 1998, Pages 205-216

doi:10.1016/S0165-3806(98)00050-9 [Cite or link using doi](#)  
 Copyright © 1998 Elsevier Science B.V. All rights reserved

**This Document**► **Abstract**

- [Full Text + Links](#)
- [PDF \(1443 K\)](#)

**Research report****Actions**

- [E-mail Article](#)

# Depolarization stimulates lamellipodia formation and axonal but not dendritic branching in cultured rat cerebral cortex neurons

G. J. A. Ramakers<sup>\*</sup>, J. Winter, T. M. Hoogland, M. B. Lequin, P. van Hulten, J. van Pelt and C. W. Pool

Netherlands Institute for Brain Research, Graduate School Neurosciences Amsterdam, Meibergdreef 33, 1105 AZ Amsterdam ZO, Netherlands

Accepted 3 March 1998. Available online 23 December 1998.

**Abstract**

Electric activity is known to have profound effects on growth cone morphology and neurite outgrowth, but the nature of the response varies strongly between neurons derived from different species or brain areas. To establish the role of electric activity in neurite outgrowth and neuronal morphogenesis of rat cerebral cortex neurons, cultured neurons were depolarized for up to 72 h and quantitatively analyzed for changes in axonal and dendritic morphology. Depolarization with 25 mM potassium chloride induced a rapid increase in lamellipodia in almost all growth cones and along both axons and dendrites. Lamellipodia formation was dependent on an influx of extracellular calcium through L-type voltage-sensitive calcium channels. Prolonged depolarization for 24 h induced an increase in total axonal length, mainly due to an increase in branching. After three days of depolarization axonal outgrowth was largely the same as in control neurons, suggesting accommodation of the growth cones to chronic depolarization. Dendrites showed very little change during the first three days in culture, and dendritic length or branching were not affected by depolarization. Thus, in early cerebral cortex neurons depolarization specifically stimulates axonal outgrowth through increased branching. This increase in branching may be a consequence of the earlier increase in lamellipodia formation. In contrast, early dendrites seem to be unable to translate the increase in lamellipodia into changes in outgrowth or

branching. This difference between axons and dendrites could be due to differences in the stabilization of the tubulin cytoskeleton.

**Author Keywords:** Neuronal morphogenesis; Growth cones; Depolarization; Neurite outgrowth and branching; Electric activity

---

\*Corresponding author. Neurons and Networks, Netherlands Institute for Brain Research, Meibergdreef 33, 1105 AZ Amsterdam ZO, The Netherlands. Fax: +31-20-6961006; E-mail: [g.ramakers@nih.knaw.nl](mailto:g.ramakers@nih.knaw.nl)

---

**Developmental Brain Research**

Volume 108, Issues 1-2 , 15 June 1998, Pages 205-216

**This Document**

► **Abstract**

- [Full Text + Links](#)
- [PDF \(1443 K\)](#)

---

**Actions**

- [E-mail Article](#)

---

[Home](#) [Journals](#) [Abstract Databases](#) [Books](#) [Reference Works](#) [My Profile](#) [Alerts](#)  [Help](#)

Send [feedback](#) to ScienceDirect

Software and compilation © 2004 ScienceDirect. All rights reserved.

ScienceDirect® is a registered trademark of Elsevier B.V.

Your use of this service is governed by [Terms and Conditions](#). Please review our [Privacy Policy](#) for details on how we protect information that you supply.

AN 1998:699079 CAPLUS

DN 130:61163

ED Entered STN: 04 Nov 1998

TI Nerve growth factor-responsive, transcription-independent outgrowth of neurites in a clonal variant of PC12 cells (PC12D)

AU Sano, Mamoru

CS Aichi Human Service Center, Institute for Developmental Research, Aichi, 480-03, Japan

SO Current Topics in Neurochemistry (1997), 1, 27-40

CODEN: CTNEFZ

PB Research Trends

DT Journal; General Review

LA English

CC 2-0 (Mammalian Hormones)

AB A **review**, with 119 refs. Most studies of the NGF-dependent outgrowth of neurites have been performed with cultured sympathetic and sensory neurons or PC12 cells. The primary neurons have had prior exposure to NGF in vivo and they require NGF for basic survival. Although PC12 cells do not require NGF for survival, the outgrowth of neurites in such cells is a consequence of their differentiation into cells that resemble sympathetic neurons in response to NGF. Thus, the outgrowth of neurites is not necessarily a direct consequence of exposure to NGF. PC12D cells, a stable variant subcloned from native PC12 cell populations, produce neurites in a rapid transcription- and translation-independent manner upon exposure to NGF, bFGF, dbcAMP or staurosporine. The rapid sprouting of neurites occurs within minutes in local regions of PC12D cells that are exposed to NGF. Recent studies in conventional PC12 cells demonstrated the close relationship between the activation of MAP kinases (ERKs) and the outgrowth of neurites in response to various agents. Simultaneous activation and rapid **nuclear** translocation of MAP kinases were also observed in PC12D cells treated with NGF or bFGF. But the activation of MAP kinases was not observed in the outgrowth of neurites induced by dbcAMP or staurosporine. In this cell line, the NGF-dependent outgrowth of neurites was not blocked by the inhibition of the activation of MAP kinases by a MEK inhibitor, PD-98059. These results indicate that the activation of MAP kinases and subsequent expression of specific genes are required for the NGF-induced differentiation of PC12 cells but this pathway is not required for the NGF-dependent outgrowth of neurites. These processes have been investigated together in most studies of conventional PC12 cells. PC12D cells provide a unique exptl. system for studies of the cellular mechanism of the NGF-induced sprouting and elongation of neurites, sep. from the transcription-dependent differentiation of the cells.

ST **review** NGF transcription neurite outgrowth

PC12 neuron signaling differentiation

IT Animal cell line

(PC12; nerve growth factor-responsive and transcription-independent outgrowth of neurites in clonal variant of PC12 cells)

IT Nerve

(differentiation; nerve growth factor-responsive and transcription-independent outgrowth of neurites in clonal variant of PC12 cells)

IT Signal transduction, biological

Transcription, genetic

(nerve growth factor-responsive and transcription-independent outgrowth of neurites in clonal variant of PC12 cells)

IT Cell differentiation

(neuronal; nerve growth factor-responsive and transcription-independent outgrowth of neurites in clonal variant of PC12 cells)

IT Axon

(outgrowth; nerve growth factor-responsive and transcription-

AN 1998:699079 CAPLUS

DN 130:61163

ED Entered STN: 04 Nov 1998

TI Nerve growth factor-responsive, transcription-independent outgrowth of neurites in a clonal variant of PC12 cells (PC12D)

AU Sano, Mamoru

CS Aichi Human Service Center, Institute for Developmental Research, Aichi, 480-03, Japan

SO Current Topics in Neurochemistry (1997), 1, 27-40

CODEN: CTNEFZ

PB Research Trends

DT Journal; General Review

LA English

CC 2-0 (Mammalian Hormones)

AB A **review**, with 119 refs. Most studies of the NGF-dependent outgrowth of neurites have been performed with cultured sympathetic and sensory neurons or PC12 cells. The primary neurons have had prior exposure to NGF in vivo and they require NGF for basic survival. Although PC12 cells do not require NGF for survival, the outgrowth of neurites in such cells is a consequence of their differentiation into cells that resemble sympathetic neurons in response to NGF. Thus, the outgrowth of neurites is not necessarily a direct consequence of exposure to NGF. PC12D cells, a stable variant subcloned from native PC12 cell populations, produce neurites in a rapid transcription- and translation-independent manner upon exposure to NGF, bFGF, dbcAMP or staurosporine. The rapid sprouting of neurites occurs within minutes in local regions of PC12D cells that are exposed to NGF. Recent studies in conventional PC12 cells demonstrated the close relationship between the activation of MAP kinases (ERKs) and the outgrowth of neurites in response to various agents. Simultaneous activation and rapid **nuclear** translocation of MAP kinases were also observed in PC12D cells treated with NGF or bFGF. But the activation of MAP kinases was not observed in the outgrowth of neurites induced by dbcAMP or staurosporine. In this cell line, the NGF-dependent outgrowth of neurites was not blocked by the inhibition of the activation of MAP kinases by a MEK inhibitor, PD-98059. These results indicate that the activation of MAP kinases and subsequent expression of specific genes are required for the NGF-induced differentiation of PC12 cells but this pathway is not required for the NGF-dependent outgrowth of neurites. These processes have been investigated together in most studies of conventional PC12 cells. PC12D cells provide a unique exptl. system for studies of the cellular mechanism of the NGF-induced sprouting and elongation of neurites, sep. from the transcription-dependent differentiation of the cells.

ST **review** NGF transcription neurite outgrowth  
PC12 neuron signaling differentiation

IT Animal cell line

(PC12; nerve growth factor-responsive and transcription-independent outgrowth of neurites in clonal variant of PC12 cells)

IT Nerve

(differentiation; nerve growth factor-responsive and transcription-independent outgrowth of neurites in clonal variant of PC12 cells)

IT Signal transduction, biological

Transcription, genetic

(nerve growth factor-responsive and transcription-independent outgrowth of neurites in clonal variant of PC12 cells)

IT Cell differentiation

(neuronal; nerve growth factor-responsive and transcription-independent outgrowth of neurites in clonal variant of PC12 cells)

IT Axon

(outgrowth; nerve growth factor-responsive and transcription-

independent outgrowth of neurites in clonal variant of PC12 cells)  
 IT 142243-02-5, Extracellular signal-regulated kinase  
 RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
 (nerve growth factor-responsive and transcription-independent outgrowth of neurites in clonal variant of PC12 cells)  
 IT 9061-61-4, Nerve growth factor  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
 (nerve growth factor-responsive and transcription-independent outgrowth of neurites in clonal variant of PC12 cells)

RE.CNT 119 THERE ARE 119 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 RE

- (1) Alessi, D; J Biol Chem 1995, V270, P27489 CAPLUS
- (2) Aletta, J; J Cell Biol 1988, V106, P1573 CAPLUS
- (3) Anderson, D; Cell 1985, V42, P649 CAPLUS
- (4) Bar-Sagi, D; Cell 1985, V42, P841 CAPLUS
- (5) Bary, D; Proc Natl Acad Sci USA 1978, V75, P5226
- (6) Berg, M; J Biol Chem 1992, V267, P13 CAPLUS
- (7) Bixby, J; J Cell Biol 1990, V111, P2725 CAPLUS
- (8) Blenis, J; Proc Natl Acad Sci USA 1993, V90, P5889 CAPLUS
- (9) Boulton, T; Cell 1991, V65, P663 CAPLUS
- (10) Burnham, P; J Neurobiol 1974, V4, P57 CAPLUS
- (11) Burstein, D; Proc Natl Acad Sci USA 1978, V75, P6059 CAPLUS
- (12) Campenot, R; Dev Biol 1982, V93, P1 MEDLINE
- (13) Chao, M; Neuron 1992, V9, P583 CAPLUS
- (14) Chen, R; Mol Cell Biol 1992, V12, P915 CAPLUS
- (15) Cobb, M; J Biol Chem 1995, V270, P14843 CAPLUS
- (16) Connolly, J; J Cell Biol 1984, V98, P457 CAPLUS
- (17) Costello, B; J Neurosci 1990, V10, P1398 CAPLUS
- (18) Cowley, S; Cell 1994, V77, P841 CAPLUS
- (19) Cox, M; Exp Cell Res 1991, V195, P423 CAPLUS
- (20) Crews, C; Cell 1993, V74, P215 CAPLUS
- (21) Davis, R; J Biol Chem 1993, V268, P14553 CAPLUS
- (22) Dichter, M; Nature 1977, V268, P501 CAPLUS
- (23) Doherty, P; Current Opinion in Cell Biol 1989, V1, P1102 MEDLINE
- (24) Doherty, P; J Cell Biol 1988, V107, P333 CAPLUS
- (25) Doherty, P; J Neurochem 1987, V49, P610 CAPLUS
- (26) Drubin, D; J Cell Biol 1985, V101, P1799 CAPLUS
- (27) Drubin, D; J Cell Biol 1988, V106, P1583 CAPLUS
- (28) Dudley, D; Proc Natl Acad Sci USA 1995, V92, P7686 CAPLUS
- (29) Erickson, A; J Biol Chem 1990, V265, P19728 CAPLUS
- (30) Friedlander, D; J Cell Biol 1986, V102, P413 CAPLUS
- (31) Frodin, M; J Biol Chem 1994, V269, P6207 CAPLUS
- (32) Fukuda, M; Oncogene 1995, V11, P239 CAPLUS
- (33) Gotoh, Y; Eur J Biochem 1990, V193, P661 CAPLUS
- (34) Greenberg, M; J Biol Chem 1985, V260, P14101 CAPLUS
- (35) Greene, L; Annual Review of Neuroscience 1980, V3, P353 CAPLUS
- (36) Greene, L; Develop Biol 1982, V91, P305 CAPLUS
- (37) Greene, L; Nature 1977, V268, P349 CAPLUS
- (38) Greene, L; Nature 1978, V276, P191 CAPLUS
- (39) Greene, L; Proc Natl Acad Sci USA 1976, V73, P2424 CAPLUS
- (40) Guerrero, I; J Cell Physiol 1986, V129, P71 CAPLUS
- (41) Gundersen, R; J Cell Biol 1980, V87, P546 CAPLUS
- (42) Hagag, N; Nature 1986, V319, P680 CAPLUS
- (43) Halbach, F; Eur J Neurosci 1992, V4, P896
- (44) Hall, D; J Cell Biol 1987, V104, P623 CAPLUS
- (45) Hanemaaijer, R; J Neurosci Res 1991, V30, P163 CAPLUS
- (46) Hashimoto, S; J Neurochem 1989, V53, P1675 CAPLUS
- (47) Heidemann, S; J Cell Biol 1985, V100, P916 CAPLUS
- (48) Hill, C; Cell 1995, V80, P199 CAPLUS

independent outgrowth of neurites in clonal variant of PC12 cells)  
 IT 142243-02-5, Extracellular signal-regulated kinase  
 RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
 (nerve growth factor-responsive and transcription-independent outgrowth of neurites in clonal variant of PC12 cells)  
 IT 9061-61-4, Nerve growth factor  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
 (nerve growth factor-responsive and transcription-independent outgrowth of neurites in clonal variant of PC12 cells)

RE.CNT 119 THERE ARE 119 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 RE

- (1) Alessi, D; J Biol Chem 1995, V270, P27489 CAPLUS
- (2) Aletta, J; J Cell Biol 1988, V106, P1573 CAPLUS
- (3) Anderson, D; Cell 1985, V42, P649 CAPLUS
- (4) Bar-Sagi, D; Cell 1985, V42, P841 CAPLUS
- (5) Bary, D; Proc Natl Acad Sci USA 1978, V75, P5226
- (6) Berg, M; J Biol Chem 1992, V267, P13 CAPLUS
- (7) Bixby, J; J Cell Biol 1990, V111, P2725 CAPLUS
- (8) Blenis, J; Proc Natl Acad Sci USA 1993, V90, P5889 CAPLUS
- (9) Boulton, T; Cell 1991, V65, P663 CAPLUS
- (10) Burnham, P; J Neurobiol 1974, V4, P57 CAPLUS
- (11) Burstein, D; Proc Natl Acad Sci USA 1978, V75, P6059 CAPLUS
- (12) Campenot, R; Dev Biol 1982, V93, P1 MEDLINE
- (13) Chao, M; Neuron 1992, V9, P583 CAPLUS
- (14) Chen, R; Mol Cell Biol 1992, V12, P915 CAPLUS
- (15) Cobb, M; J Biol Chem 1995, V270, P14843 CAPLUS
- (16) Connolly, J; J Cell Biol 1984, V98, P457 CAPLUS
- (17) Costello, B; J Neurosci 1990, V10, P1398 CAPLUS
- (18) Cowley, S; Cell 1994, V77, P841 CAPLUS
- (19) Cox, M; Exp Cell Res 1991, V195, P423 CAPLUS
- (20) Crews, C; Cell 1993, V74, P215 CAPLUS
- (21) Davis, R; J Biol Chem 1993, V268, P14553 CAPLUS
- (22) Dichter, M; Nature 1977, V268, P501 CAPLUS
- (23) Doherty, P; Current Opinion in Cell Biol 1989, V1, P1102 MEDLINE
- (24) Doherty, P; J Cell Biol 1988, V107, P333 CAPLUS
- (25) Doherty, P; J Neurochem 1987, V49, P610 CAPLUS
- (26) Drubin, D; J Cell Biol 1985, V101, P1799 CAPLUS
- (27) Drubin, D; J Cell Biol 1988, V106, P1583 CAPLUS
- (28) Dudley, D; Proc Natl Acad Sci USA 1995, V92, P7686 CAPLUS
- (29) Erickson, A; J Biol Chem 1990, V265, P19728 CAPLUS
- (30) Friedlander, D; J Cell Biol 1986, V102, P413 CAPLUS
- (31) Frodin, M; J Biol Chem 1994, V269, P6207 CAPLUS
- (32) Fukuda, M; Oncogene 1995, V11, P239 CAPLUS
- (33) Gotoh, Y; Eur J Biochem 1990, V193, P661 CAPLUS
- (34) Greenberg, M; J Biol Chem 1985, V260, P14101 CAPLUS
- (35) Greene, L; Annual Review of Neuroscience 1980, V3, P353 CAPLUS
- (36) Greene, L; Develop Biol 1982, V91, P305 CAPLUS
- (37) Greene, L; Nature 1977, V268, P349 CAPLUS
- (38) Greene, L; Nature 1978, V276, P191 CAPLUS
- (39) Greene, L; Proc Natl Acad Sci USA 1976, V73, P2424 CAPLUS
- (40) Guerrero, I; J Cell Physiol 1986, V129, P71 CAPLUS
- (41) Gundersen, R; J Cell Biol 1980, V87, P546 CAPLUS
- (42) Hagag, N; Nature 1986, V319, P680 CAPLUS
- (43) Halbach, F; Eur J Neurosci 1992, V4, P896
- (44) Hall, D; J Cell Biol 1987, V104, P623 CAPLUS
- (45) Hanemaaijer, R; J Neurosci Res 1991, V30, P163 CAPLUS
- (46) Hashimoto, S; J Neurochem 1989, V53, P1675 CAPLUS
- (47) Heidemann, S; J Cell Biol 1985, V100, P916 CAPLUS
- (48) Hill, C; Cell 1995, V80, P199 CAPLUS

- (49) Hoshi, H; J Biol Chem 1988, V263, P5396
- (50) Iwasaki, S; J Biol Chem 1996, V271, P17360 CAPLUS
- (51) Joneson, T; Science 1996, V271, P810 CAPLUS
- (52) Katoh-Semba, R; J Neurosci Res 1987, V17, P36 CAPLUS
- (53) Klinz, F; J neurosci Res 1996, V46, P720 CAPLUS
- (54) Kongsamut, S; Proc Natl Acad Sci USA 1986, V83, P2243 CAPLUS
- (55) Lamoureux, P; J Cell Biol 1990, V110, P71 CAPLUS
- (56) Landreth, G; J Cell Biol 1985, V100, P677 CAPLUS
- (57) Lemmon, V; J Neurosci 1992, V12, P818 CAPLUS
- (58) Leonard, D; J Cell Biol 1988, V106, P181 CAPLUS
- (59) Levi, A; Science 1985, V229, P393 CAPLUS
- (60) Levi-Montalcini, R; EMBO J 1987, V6, P1145 CAPLUS
- (61) Lindenbaum, M; J Biol Chem 1987, V262, P605 CAPLUS
- (62) Lloyd, E; J Neurochem 1992, V59, P1099 CAPLUS
- (63) Machida, C; Neuron 1989, V2, P1587 CAPLUS
- (64) Mandel, G; Proc Natl Acad Sci USA 1988, V85, P924 CAPLUS
- (65) Mann, D; J Neurochem 1989, V53, P1581 CAPLUS
- (66) Margolis, R; Biochemistry 1986, V25, P3463 CAPLUS
- (67) Marshall, C; Cell 1995, V80, P179 CAPLUS
- (68) Milbrant, J; Science 1987, V238, P797
- (69) Miyasaka, T; J Biol Chem 1990, V265, P4730 CAPLUS
- (70) Miyasaka, T; Proc Natl Acad Sci USA 1991, V88, P2653 CAPLUS
- (71) Nakafuku, M; J Biol Chem 1992, V267, P19448 CAPLUS
- (72) Nishiki, T; Biochem Biophys Res Comm 1990, V167, P265 CAPLUS
- (73) Nobes, C; Cell 1995, V81, P53 CAPLUS
- (74) Noda, M; Nature 1985, V318, P73 CAPLUS
- (75) Obermeier, A; EMBO J 1994, V13, P1585 CAPLUS
- (76) Obermeier, A; J Biol Chem 1993, P22963 CAPLUS
- (77) Okumura, N; J Biochem (Tokyo) 1994, V115, P304 CAPLUS
- (78) O'Laque, P; Proc Natl Acad Sci USA 1980, V77, P1701 CAPLUS
- (79) Pang, L; J Biol Chem 1995, V270, P13585 CAPLUS
- (80) Partlow, L; J Neurochem 1971, V18, P2101 CAPLUS
- (81) Paves, H; Exp Cell Res 1990, V186, P218 CAPLUS
- (82) Pelech, S; Science 1992, V257, P1355 MEDLINE
- (83) Peng, X; Neuron 1995, V15, P395 CAPLUS
- (84) Prentice, H; EMBO J 1987, V6, P1859 CAPLUS
- (85) Pulverer, B; Nature 1991, V353, P670 CAPLUS
- (86) Qiu, M; Neuron 1992, V9, P705 CAPLUS
- (87) Rasouly, D; Mol Pharmacol 1992, V42, P35 CAPLUS
- (88) Ray, L; J Biol Chem V263, P12721 CAPLUS
- (89) Ridley, A; Cell 1992, V70, P389 CAPLUS
- (90) Ridley, A; Cell 1992, V70, P401 CAPLUS
- (91) Sano, M; Brain Res 1988, V459, P404 CAPLUS
- (92) Sano, M; Brain Res 1994, V639, P115 CAPLUS
- (93) Sano, M; Brain Res 1994, V656, P210 CAPLUS
- (94) Sano, M; Brain Res 1995, V688, P213 CAPLUS
- (95) Sano, M; Brain Res 1996, V742, P195 CAPLUS
- (96) Sano, M; Brain Res 1996, V730, P212 CAPLUS
- (97) Sano, M; Cell Str Func 1992, V17, P341 CAPLUS
- (98) Sano, M; J Neurochem 1990, V55, P427 CAPLUS
- (99) Sano, M; J Neurochem 1992, V58, P837 CAPLUS
- (100) Schubert, D; Proc Natl Acad Sci USA 1977, V74, P2579 CAPLUS
- (101) Schuch, U; Neuron 1989, V3, P13 CAPLUS
- (102) Stephens, R; Neuron 1994, V12, P691 CAPLUS
- (103) Szeberenyi, J; Mol Cell Biol 1990, V10, P5324 CAPLUS
- (104) Takahashi, M; Brain Res 1985, V341, P381 CAPLUS
- (105) Teng, K; J Biol Chem 1995, V270, P20677 CAPLUS
- (106) Tigyi, G; J Neurochem 1996, V66, P537 CAPLUS
- (107) Tirone, F; Proc Natl Acad Sci USA 1989, V86, P2088 CAPLUS
- (108) Tischler, A; J Neurosci 1986, V17, P1719
- (109) Traverse, S; Biochem J 1992, V288, P351 CAPLUS
- (110) Tsao, H; J Biol Chem 1990, V265, P15471 CAPLUS



- (49) Hoshi, H; J Biol Chem 1988, V263, P5396
- (50) Iwasaki, S; J Biol Chem 1996, V271, P17360 CAPLUS
- (51) Joneson, T; Science 1996, V271, P810 CAPLUS
- (52) Katoh-Semba, R; J Neurosci Res 1987, V17, P36 CAPLUS
- (53) Klinz, F; J neurosci Res 1996, V46, P720 CAPLUS
- (54) Kongsamut, S; Proc Natl Acad Sci USA 1986, V83, P2243 CAPLUS
- (55) Lamoureux, P; J Cell Biol 1990, V110, P71 CAPLUS
- (56) Landreth, G; J Cell Biol 1985, V100, P677 CAPLUS
- (57) Lemmon, V; J Neurosci 1992, V12, P818 CAPLUS
- (58) Leonard, D; J Cell Biol 1988, V106, P181 CAPLUS
- (59) Levi, A; Science 1985, V229, P393 CAPLUS
- (60) Levi-Montalcini, R; EMBO J 1987, V6, P1145 CAPLUS
- (61) Lindenbaum, M; J Biol Chem 1987, V262, P605 CAPLUS
- (62) Lloyd, E; J Neurochem 1992, V59, P1099 CAPLUS
- (63) Machida, C; Neuron 1989, V2, P1587 CAPLUS
- (64) Mandel, G; Proc Natl Acad Sci USA 1988, V85, P924 CAPLUS
- (65) Mann, D; J Neurochem 1989, V53, P1581 CAPLUS
- (66) Margolis, R; Biochemistry 1986, V25, P3463 CAPLUS
- (67) Marshall, C; Cell 1995, V80, P179 CAPLUS
- (68) Milbrant, J; Science 1987, V238, P797
- (69) Miyasaka, T; J Biol Chem 1990, V265, P4730 CAPLUS
- (70) Miyasaka, T; Proc Natl Acad Sci USA 1991, V88, P2653 CAPLUS
- (71) Nakafuku, M; J Biol Chem 1992, V267, P19448 CAPLUS
- (72) Nishiki, T; Biochem Biophys Res Comm 1990, V167, P265 CAPLUS
- (73) Nobes, C; Cell 1995, V81, P53 CAPLUS
- (74) Noda, M; Nature 1985, V318, P73 CAPLUS
- (75) Obermeier, A; EMBO J 1994, V13, P1585 CAPLUS
- (76) Obermeier, A; J Biol Chem 1993, P22963 CAPLUS
- (77) Okumura, N; J Biochem (Tokyo) 1994, V115, P304 CAPLUS
- (78) O'Lague, P; Proc Natl Acad Sci USA 1980, V77, P1701 CAPLUS
- (79) Pang, L; J Biol Chem 1995, V270, P13585 CAPLUS
- (80) Partlow, L; J Neurochem 1971, V18, P2101 CAPLUS
- (81) Paves, H; Exp Cell Res 1990, V186, P218 CAPLUS
- (82) Pelech, S; Science 1992, V257, P1355 MEDLINE
- (83) Peng, X; Neuron 1995, V15, P395 CAPLUS
- (84) Prentice, H; EMBO J 1987, V6, P1859 CAPLUS
- (85) Pulverer, B; Nature 1991, V353, P670 CAPLUS
- (86) Qiu, M; Neuron 1992, V9, P705 CAPLUS
- (87) Rasouly, D; Mol Pharmacol 1992, V42, P35 CAPLUS
- (88) Ray, L; J Biol Chem V263, P12721 CAPLUS
- (89) Ridley, A; Cell 1992, V70, P389 CAPLUS
- (90) Ridley, A; Cell 1992, V70, P401 CAPLUS
- (91) Sano, M; Brain Res 1988, V459, P404 CAPLUS
- (92) Sano, M; Brain Res 1994, V639, P115 CAPLUS
- (93) Sano, M; Brain Res 1994, V656, P210 CAPLUS
- (94) Sano, M; Brain Res 1995, V688, P213 CAPLUS
- (95) Sano, M; Brain Res 1996, V742, P195 CAPLUS
- (96) Sano, M; Brain Res 1996, V730, P212 CAPLUS
- (97) Sano, M; Cell Str Func 1992, V17, P341 CAPLUS
- (98) Sano, M; J Neurochem 1990, V55, P427 CAPLUS
- (99) Sano, M; J Neurochem 1992, V58, P837 CAPLUS
- (100) Schubert, D; Proc Natl Acad Sci USA 1977, V74, P2579 CAPLUS
- (101) Schuch, U; Neuron 1989, V3, P13 CAPLUS
- (102) Stephens, R; Neuron 1994, V12, P691 CAPLUS
- (103) Szeberenyi, J; Mol Cell Biol 1990, V10, P5324 CAPLUS
- (104) Takahashi, M; Brain Res 1985, V341, P381 CAPLUS
- (105) Teng, K; J Biol Chem 1995, V270, P20677 CAPLUS
- (106) Tigyi, G; J Neurochem 1996, V66, P537 CAPLUS
- (107) Tirone, F; Proc Natl Acad Sci USA 1989, V86, P2088 CAPLUS
- (108) Tischler, A; J Neurosci 1986, V17, P1719
- (109) Traverse, S; Biochem J 1992, V288, P351 CAPLUS
- (110) Tsao, H; J Biol Chem 1990, V265, P15471 CAPLUS

- (111) Twiss, J; J Neurochem 1995, V64, P550 CAPLUS
- (112) Vaillancourt, R; Mol Cell Biol 1995, V15, P3644 CAPLUS
- (113) Virdee, K; J Neurochem 1996, V67, P1801 CAPLUS
- (114) Williams, E; J Cell Biol 1994, V124, P1029 CAPLUS
- (115) Wood, K; Cell 1992, V68, P1041 CAPLUS
- (116) Wood, K; Proc Natl Acad Sci USA 1993, V90, P5016 CAPLUS
- (117) Wu, B; J Biol Chem 1989, V264, P9000 CAPLUS
- (118) Young, S; FEBS Lett 1994, V338, P212 CAPLUS
- (119) Zhang, Z; J Biol Chem 1993, V268, P5557 CAPLUS

- (111) Twiss, J; J Neurochem 1995, V64, P550 CAPLUS
- (112) Vaillancourt, R; Mol Cell Biol 1995, V15, P3644 CAPLUS
- (113) Virdee, K; J Neurochem 1996, V67, P1801 CAPLUS
- (114) Williams, E; J Cell Biol 1994, V124, P1029 CAPLUS
- (115) Wood, K; Cell 1992, V68, P1041 CAPLUS
- (116) Wood, K; Proc Natl Acad Sci USA 1993, V90, P5016 CAPLUS
- (117) Wu, B; J Biol Chem 1989, V264, P9000 CAPLUS
- (118) Young, S; FEBS Lett 1994, V338, P212 CAPLUS
- (119) Zhang, Z; J Biol Chem 1993, V268, P5557 CAPLUS

ANSWER 6 OF 6 MEDLINE on STN  
AN 96162646 MEDLINE  
DN PubMed ID: 8581313  
TI Multiple factors govern intraretinal axon guidance: a time-lapse study.  
AU Brittis P A; Silver J  
CS Department of Neurosciences, Case Western Reserve University, Cleveland, Ohio 44106, USA.  
SO Molecular and cellular neurosciences, (1995 Oct) 6 (5) 413-32.  
Journal code: 9100095. ISSN: 1044-7431.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199603  
ED Entered STN: 19960327  
Last Updated on STN: 19960327  
Entered Medline: 19960319  
AB In this study, the multiple factors that govern the unidirectional path of intraretinal axons, as well as the cellular movements prior to and during early axonogenesis, were investigated using time-lapse videomicroscopy. For several hours prior to overt axon elongation, young retinal ganglion cells send out transient minor processes in all directions at the pial surface. Time-lapse analysis of the chondroitin sulfate (CS)-containing matrix that has been suggested to play an important role in regulating this early differentiative event revealed the dynamic, wavelike properties of this extracellular matrix component. As the CS matrix dissipates across the immature ganglion cells, only one minor process, away from the highest concentration of CS peripherally and in the direction of the optic fissure centrally, is retained and becomes the mature axon. Focal concentrations of Ll appear at points of **neurite** contact with previously established axons, suggesting that this growth-promoting molecule is also involved with establishing the precise, unidirectional **outgrowth** pattern of retinal ganglion cell axons. NCAM was diffusely distributed on neural elements and on the neuroepithelial endfeet in the central and peripheral retina and, thus, may not be an essential unidirectional axon growth cue. Growth cones mechanically deflected 180 degrees from the optic fissure after the CS wave had receded from the central retina had morphologies and rates of elongation similar to those oriented in the proper direction. Growth cones deflected obliquely toward the ventral retinal periphery entered a territory of increasing CS-containing proteoglycan matrix and neurons with minor processes. As these deflected axons entered more deeply into this region they slowed down and sent out long transient branchlike processes. These observations illustrate the complex organization of the changing cell surface and matrix components within the retina during axonogenesis and axon **outgrowth**. The results also elucidate the potential importance of a cell state where immature neurons probe their environment via minor processes. These specialized **neurites** may provide the neuron with a way to sample a full 360 degrees of terrain around them. This method of exploring the environment could afford the cell a mechanism with which to sample, summate, and respond to physical structures as well as simultaneously occurring negative and positive molecular influences that are distributed unequally on either side of the **cell body**.  
CT Check Tags: Female; Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S. Animals  
Axons: ME, metabolism  
\*Axons: PH, physiology  
Axons: UL, ultrastructure  
Image Processing, Computer-Assisted  
Immunohistochemistry  
Microscopy, Electron

ANSWER 6 OF 6 MEDLINE on STN  
AN 96162646 MEDLINE  
DN PubMed ID: 8581313  
TI Multiple factors govern intraretinal axon guidance: a time-lapse study.  
AU Brittis P A; Silver J  
CS Department of Neurosciences, Case Western Reserve University, Cleveland, Ohio 44106, USA.  
SO Molecular and cellular neurosciences, (1995 Oct) 6 (5) 413-32.  
Journal code: 9100095. ISSN: 1044-7431.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199603  
ED Entered STN: 19960327  
Last Updated on STN: 19960327  
Entered Medline: 19960319  
AB In this study, the multiple factors that govern the unidirectional path of intraretinal axons, as well as the cellular movements prior to and during early axonogenesis, were investigated using time-lapse videomicroscopy. For several hours prior to overt axon elongation, young retinal ganglion cells send out transient minor processes in all directions at the pial surface. Time-lapse analysis of the chondroitin sulfate (CS)-containing matrix that has been suggested to play an important role in regulating this early differentiative event revealed the dynamic, wavelike properties of this extracellular matrix component. As the CS matrix dissipates across the immature ganglion cells, only one minor process, away from the highest concentration of CS peripherally and in the direction of the optic fissure centrally, is retained and becomes the mature axon. Focal concentrations of L1 appear at points of **neurite** contact with previously established axons, suggesting that this growth-promoting molecule is also involved with establishing the precise, unidirectional **outgrowth** pattern of retinal ganglion cell axons. NCAM was diffusely distributed on neural elements and on the neuroepithelial endfeet in the central and peripheral retina and, thus, may not be an essential unidirectional axon growth cue. Growth cones mechanically deflected 180 degrees from the optic fissure after the CS wave had receded from the central retina had morphologies and rates of elongation similar to those oriented in the proper direction. Growth cones deflected obliquely toward the ventral retinal periphery entered a territory of increasing CS-containing proteoglycan matrix and neurons with minor processes. As these deflected axons entered more deeply into this region they slowed down and sent out long transient branchlike processes. These observations illustrate the complex organization of the changing cell surface and matrix components within the retina during axonogenesis and axon **outgrowth**. The results also elucidate the potential importance of a cell state where immature neurons probe their environment via minor processes. These specialized **neurites** may provide the neuron with a way to sample a full 360 degrees of terrain around them. This method of exploring the environment could afford the cell a mechanism with which to sample, summate, and respond to physical structures as well as simultaneously occurring negative and positive molecular influences that are distributed unequally on either side of the **cell body**.  
CT Check Tags: Female; Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S. Animals  
Axons: ME, metabolism  
\*Axons: PH, physiology  
Axons: UL, ultrastructure  
Image Processing, Computer-Assisted  
Immunohistochemistry  
Microscopy, Electron

Rats  
Rats, Sprague-Dawley  
Retinal Ganglion Cells: ME, metabolism  
\*Retinal Ganglion Cells: PH, physiology  
Time Factors

=>

Rats  
Rats, Sprague-Dawley  
Retinal Ganglion Cells: ME, metabolism  
\*Retinal Ganglion Cells: PH, physiology  
Time Factors

=>

ANSWER 4 OF 6 CAPLUS COPYRIGHT 2004 ACS on STN

AN 1985:520824 CAPLUS

DN 103:120824

ED Entered STN: 19 Oct 1985

TI The effect of exogenous gangliosides on neurons in culture: a morphometric analysis

AU Massarelli, R.; Ferret, B.; Gorio, A.; Durand, M.; Dreyfus, H.

CS Cent. Neurochim., CNRS, Strasbourg, 67084, Fr.

SO International Journal of Developmental Neuroscience (1985), 3(4), 341-8

CODEN: IJDND6; ISSN: 0736-5748

DT Journal

LA English

CC 13-6 (Mammalian Biochemistry)

AB Cultures of isolated neurons were treated with a purified preparation of gangliosides (10-5M and 10-9M) added to the cell growth medium at the 3rd day in culture, and a morphometric anal. of the cells was performed with an **image** analyzer after 1 and 4 days of treatment. The number of cells and the area of the **cell bodies** were increased following the treatment. Also, there was apparently a sprouting effect of the glycolipids on the number of secondary neuronal processes and an increase in the length of the primary **neurites**. The present data and other biochem. evidence (Dreyfus, H., et al., 1984) suggest that the addition of exogenous gangliosides may have a trophic effect on neurons, greatly enhances the number of cell-to-cell contacts, and possibly stimulates cell proliferation and differentiation.

ST neuron growth culture ganglioside; **neurite outgrowth**  
ganglioside

IT Nerve

(growth and morphol. of, in culture, gangliosides effect on)

IT Gangliosides

RL: BIOL (Biological study)

(neuron growth and morphol. and axon **outgrowth** in culture  
response to)

IT Animal tissue culture

(neuron growth and morphol. and axon sprouting in, gangliosides effect  
on)

IT Nerve

(axon, length of, in neurons in culture, gangliosides effect on)



ANSWER 4 OF 6 CAPLUS COPYRIGHT 2004 ACS on STN

AN 1985:520824 CAPLUS

DN 103:120824

ED Entered STN: 19 Oct 1985

TI The effect of exogenous gangliosides on neurons in culture: a morphometric analysis

AU Massarelli, R.; Ferret, B.; Gorio, A.; Durand, M.; Dreyfus, H.

CS Cent. Neurochim., CNRS, Strasbourg, 67084, Fr.

SO International Journal of Developmental Neuroscience (1985), 3(4), 341-8

CODEN: IJDND6; ISSN: 0736-5748

DT Journal

LA English

CC 13-6 (Mammalian Biochemistry)

AB Cultures of isolated neurons were treated with a purified preparation of gangliosides (10-5M and 10-9M) added to the cell growth medium at the 3rd day in culture, and a morphometric anal. of the cells was performed with an **image** analyzer after 1 and 4 days of treatment. The number of cells and the area of the **cell bodies** were increased following the treatment. Also, there was apparently a sprouting effect of the glycolipids on the number of secondary neuronal processes and an increase in the length of the primary **neurites**. The present data and other biochem. evidence (Dreyfus, H., et al., 1984) suggest that the addition of exogenous gangliosides may have a trophic effect on neurons, greatly enhances the number of cell-to-cell contacts, and possibly stimulates cell proliferation and differentiation.

ST neuron growth culture ganglioside; **neurite outgrowth**  
ganglioside

IT Nerve

(growth and morphol. of, in culture, gangliosides effect on)

IT Gangliosides

RL: BIOL (Biological study)

(neuron growth and morphol. and axon **outgrowth** in culture  
response to)

IT Animal tissue culture

(neuron growth and morphol. and axon sprouting in, gangliosides effect  
on)

IT Nerve

(axon, length of, in neurons in culture, gangliosides effect on)

AN 1999:198128 CAPLUS  
 DN 131:16923  
 ED Entered STN: 29 Mar 1999  
 TI Effects of different fragments of tenascin-R on neuron morphology in vitro  
 AU Xu, Hanpeng; Xiao, Huasheng; Wang, Haojun; Liang, Zhe; Gong, Ju  
 CS The Institute of Neuroscience, Fourth Military Medical University, Xi'an, 710033, Peop. Rep. China  
 SO Journal of Medical Colleges of PLA (1998), 13(4), 272-275  
 CODEN: JMCPE6; ISSN: 1000-1948  
 PB Journal of Medical Colleges of PLA, Editorial Board  
 DT Journal  
 LA English  
 CC 13-6 (Mammalian Biochemistry)  
 AB To investigate the effects of different tenascin-R fragments on morphol. changes of neurons in vitro, cell suspension were prepared from E14-15 mouse embryo spinal cords by mech. dissection and trypsin digestion. The cells were cultured in dishes coated with different bacterial expressed tenascin-R fragments. After being cultured in serum-free medium for 26 h, the cells were fixed and stained by ABC immunocytochem. method for NSE. The cell number and **neurite** length were measured by a stereol. method using an **image** anal. system, and the data was analyzed statistically. The cells grew well in the serum-free medium for 26 h. Three types of cells were identified: (1) phase-bright cells with single or double **neurites**; (2) phase-dark cells with well branching **neurites**; (3) flat cells with 2-4 round vesicles in the **cell body** and radio-like **neurites**. The cell number and the **neurites** length were influenced by different tenascin-R fragments. It was found that FN1-2 fragment inhibited **neurite outgrowth**. Different tenascin-R fragments that were used as substrate exert varying effects on cultured neural cells, adhesion or anti-adhesion of cells, promotion or inhibition of the growth of **neurites**. These influences were mediated through receptors on the cell membrane. This study may provide some clues to the search for the search for the different receptors which may play essential roles in neuronal development and plasticity.  
 ST tenascin R neuron morphol  
 IT Nerve  
     (neuron; tenascin-R fragment effect on neuron)  
 IT Axon  
     (**outgrowth**; tenascin-R fragment effect on neuron)  
 IT Cell adhesion  
     Cell morphology  
     Spinal cord  
     (tenascin-R fragment effect on neuron)  
 IT Tenascins  
     RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
     (tenascin-R; tenascin-R fragment effect on neuron)  
 RE.CNT 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 RE  
 (1) Chiquet, E; Curr Opin Cell Biol 1995, V7(5), P715  
 (2) Dorries, U; J Neurosci Res 1996, V43(4), P420 MEDLINE  
 (3) Gundersen, H; APMIS 1988, V96(5), P379 MEDLINE  
 (4) Gundersen, H; J Microsc 1987, V147(Pt3), P229  
 (5) Hay Elizabeth, D; Cell Biol of ECM 2nd ed 1991, P419  
 (6) Kruse, J; Nature 1985, V316, P146 CAPLUS  
 (7) Pesheva, P; J Cell Biol 1989, V109(4), P1765 CAPLUS  
 (8) Pesheva, P; Neuron 1992, V6(1), P69  
 (9) Sjmith, B; Cell Adv Com 1996, V3(6), P451  
 (10) Taylor, J; J Neurosci Res 1993, V35(4), P347 CAPLUS  
 (11) Vidal, L; J Neurosci Methods 1995, V56(1), P89 CAPLUS

(12) Villa, M; J Neurosci Methods 1995, V56(1), P77 MEDLINE

(12) Villa, M; J Neurosci Methods 1995, V56(1), P77 MEDLINE

d his

(FILE 'HOME' ENTERED AT 14:09:50 ON 16 MAR 2004)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, CANCERLIT, JAPIO' ENTERED AT  
14:10:15 ON 16 MAR 2004

L1 19246 S (NEURITE OUTGROWTH)  
L2 39 S L1 AND LUMINES?  
L3 30 DUPLICATE REMOVE L2 (9 DUPLICATES REMOVED)  
L4 731 S L1 AND REVIEW?  
L5 477 DUPLICATE REMOVE L4 (254 DUPLICATES REMOVED)  
L6 34 S L5 AND NUCLE?  
L7 34 S L6 NOT L3  
L8 3714 S NEURITE? AND (CELL BOD?)  
L9 22 S L8 AND LUMINESC?  
L10 7 S L9 AND IMAGE?  
L11 4 DUPLICATE REMOVE L10 (3 DUPLICATES REMOVED)  
L12 85 S L8 AND IMAGE?  
L13 39 DUPLICATE REMOVE L12 (46 DUPLICATES REMOVED)  
L14 39 S L13 NOT L11  
L15 0 S L14 AND LUMINE?  
L16 0 S L14 AND LUMINE?  
L17 6 S L14 AND OUTGROWTH?

=>

d his

(FILE 'HOME' ENTERED AT 14:09:50 ON 16 MAR 2004)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, CANCERLIT, JAPIO' ENTERED AT  
14:10:15 ON 16 MAR 2004

L1 19246 S (NEURITE OUTGROWTH)  
L2 39 S L1 AND LUMINES?  
L3 30 DUPLICATE REMOVE L2 (9 DUPLICATES REMOVED)  
L4 731 S L1 AND REVIEW?  
L5 477 DUPLICATE REMOVE L4 (254 DUPLICATES REMOVED)  
L6 34 S L5 AND NUCLE?  
L7 34 S L6 NOT L3  
L8 3714 S NEURITE? AND (CELL BOD?)  
L9 22 S L8 AND LUMINESC?  
L10 7 S L9 AND IMAGE?  
L11 4 DUPLICATE REMOVE L10 (3 DUPLICATES REMOVED)  
L12 85 S L8 AND IMAGE?  
L13 39 DUPLICATE REMOVE L12 (46 DUPLICATES REMOVED)  
L14 39 S L13 NOT L11  
L15 0 S L14 AND LUMINE?  
L16 0 S L14 AND LUMINE?  
L17 6 S L14 AND OUTGROWTH?

=>